

Analysis of Plant Remains from SBA-3587

prepared by:
Amber M. VanDerwarker
Jennifer V. Alvarado

Report Submitted to Ray Corbett, PhD.

Table of Contents

Introduction	2
Recovery and Preservation Bias	2
Laboratory Procedures	3
Basic Results	3
Table 1 (Summary of plant taxa for SBA-3587 flotation samples)	4
References Cited	4

Introduction

Archaeological plant and animal assemblages represent only a small fraction of what was originally used and deposited by humans in open-air settings. Natural and cultural factors can significantly modify organic remains, resulting in recovered assemblages that differ dramatically from the original deposits. As archaeologists, we examine collections that have undergone a series of processes—from the original selection of plants and animals by humans, to food preparation, cooking, discard, animal and insect scavenging, burial, decay, and weathering, to the recovery of food residues by archaeologists. Using standard methodological procedures for sampling, quantification, and analysis allows us to make sense of our assemblages in spite of the deleterious effects of these processes. Here we report on the identification and analysis of the archaeobotanical assemblage from one test unit ranging from 40-120 cm depth levels at the SBA-3587 Chumash site in the foothills of Carpinteria, California.

Recovery and Preservation Bias

The circumstances under which plants preserve best archaeologically involve extreme conditions (e.g., exceptionally wet, dry, or cold environments) that prohibit decomposition of organic matter (Miksicek 1987). Plants can also preserve through exposure to fire, which can transform plant material from organic matter into carbon (Miksicek 1987). The likelihood that a plant will become carbonized varies according to the type of plant, how it is prepared and used, and whether it has a dense or fragile structure (Scarry 1986). Plants that are eaten whole are less likely to produce discarded portions that may find their way into a fire. Plants that require the removal of inedible portions (e.g., hickory nutshell, corn cobs) are more likely to find their way into a fire, and thus into the archaeological record. Inedible plant parts represent intentional discard that is often burned as fuel. Moreover, because inedible portions tend to be dense and fibrous, they are more likely to survive the process of carbonization than the edible parts (e.g., hickory nutshell vs. nutmeats). Physical characteristics are also important for determining whether or not a plant will survive a fire. Thick, dense nutshells are more likely to survive a fire than smaller, more fragile grass seeds. Food preparation activities also affect potential plant carbonization. The simple process of cooking provides the opportunity for carbonization through cooking accidents. Foods that are conventionally eaten raw, however, are less likely to be deposited in fires than cooked foods.

Some plants that find their way into the archaeological record in carbonized form were not eaten at all. Wood fuel is the most obvious example. Burned house structures can also yield carbonized plant deposits, and these deposits often differ dramatically from refuse deposits (Scarry 1986). Other non-food plants that become carbonized are incidental inclusions, such as seeds blown by wind dispersal (Miksicek 1987; Minnis 1981; Scarry 1986). Indeed, most secondary invaders are weedy species with lots of seeds (e.g., cheno/am plants) (Minnis 1981).

While we cannot ever hope to know the absolute quantities or importance of different plants in any past subsistence economy, the preservation and recovery biases discussed above do not prohibit quantitative analyses of archaeobotanical assemblages. The most commonly used plant resources in any subsistence economy are more likely to be subject to activities that result in carbonization (e.g., through fuel use and accidental burning) and ultimately, deposition (Scarry 1986; Yarnell 1982). Thus, we can quantitatively examine the relative importance of commonly used plant resources through time and across space.

Laboratory Procedures

Eight flotation samples from SBA-3587 were collected with variable volumes, ranging from 8 to 12 liters. Both the light and heavy fractions of the flotation samples were analyzed. Although the materials from the light and heavy fractions were processed and sorted separately, data from the two fractions were combined for analysis. According to standard practice, the light fractions were weighed and then sifted through 2.0 mm, 1.0mm, and 0.7 mm standard geological sieves. The heavy fraction component of each sample was weighed and then sifted through 2.0 mm and 1.4 mm standard geological sieves. Carbonized plant remains from both fractions were sorted in entirety down to the first two sieve sizes for each respective fraction with the aid of a stereoscopic microscope (10–40 X). Residue less than 1.0 mm in size (light fraction) and 1.4 mm in size (heavy fraction) was scanned for seeds, which were removed and counted.

Botanical materials were identified with reference to the paleoethnobotanical comparative collection at the University of California, Santa Barbara (UCSB) paleoethnobotany lab, various seed identification manuals (Martin and Barkley 1961; Delorit 1970), the USDA pictorial website (<http://www.ars-grin.gov/npgs/images/sbml/>), Timbrook (2007) and Minnis (2004) which allowed us to identify the range of taxa native to the region. All plant specimens were identified to the lowest possible taxonomic level. Taxonomic identification was not always possible—some plant specimens lacked diagnostic features altogether or were too highly fragmented. As a result, these specimens were classified as “unidentified” or “unidentified seed.” In other cases, probable identifications were made—for example, if a specimen closely resembled acorn, but a clear taxonomic distinction was not possible (e.g., the specimen was highly fragmented), then the specimen was identified as a probable acorn and recorded as “acorn cf.”.

Once the plant specimens were sorted and identified, we recorded counts, weights (in grams), and provenience information. Wood was weighed but not counted, and no wood identification was conducted. Generally, most of the seeds identified in the samples were too small to weigh, and thus only counts were recorded. Other than counts and weights, no other measurements were taken on any specimens. We also sub-sampled selected samples that were extremely large. These samples were weighed and then systematically split using a riffle splitter; some samples were split in half and others in quarters depending on the overall weight of the sample. Counts and weights from the selected subsample were extrapolated using the total sample weight.

Basic Results

This section presents the results of the identification of the carbonized plant remains from SBA-3587. Table 1 lists the common and taxonomic names of all identified species. Raw counts and weights are provided for each taxon; plant weight and wood weight are also provided. Combined, these samples yielded 10 plant taxa. Possible acorn (*Quercus sp.*) nutshell was identified. Acorns were gathered from the trees and the ground, then dried and shelled. Because of the tannins present in acorn nutmeat, leaching the tannins was necessary. This could be done by placing ground acorn meal in a bowl or leaching basin, which could then be filled with water to soak the meal and reduce the acidity of the acorns. After the leaching process, acorn meal could be incorporated into acorn mush and possibly breads (Timbrook 2007). Possible California bay (*Umbellularia californica*) was also identified. Bay leaves were used medicinally to help treat colds and headaches. It is unknown whether the Chumash also utilized the fruit of this plant (Timbrook 2007). Wild cucumber (*Marah macrocarpus*) rind overwhelmingly dominates the plant assemblage; the seeds from this plant have medicinal uses. The roasted seeds also could have been combined with other plants to create paint pigment (Timbrook 2007).

Amaranth (*Amaranthus. sp.*) and Chenopodium (*Chenopodium sp.*) seeds identified in the samples could have been eaten or ground into a meal. The leaves of both species also could have been eaten. One fruit taxon, nightshade (*Solanum sp.*) was identified. The berries could have been eaten raw or boiled. A possible seed from the legume family (*Fabaceae*) could represent clover, which has edible leaves and shoots (Timbrook 2007). Other seeds that probably represent incidental inclusions in the assemblage include St. johnswort (*Hypericum sp.*), violet (*Viola sp.*) and canarygrass (*Phalaris sp.*).

Table 1. Summary of plant taxa from SBA-3587

N of Samples		8	
Plant Weight (grams)		1.63	
Wood Weight (grams)		0.68	
<u>Common Name</u>	<u>Taxonomic Name</u>	<u>Count (n)</u>	<u>Weight (g)</u>
Acorn cf.	<i>Quercus sp.</i>	5	0.03
Amaranth	<i>Amaranthus sp.</i>	1	0
	<i>Umbellularia</i>		
California Bay, cf.	<i>californica</i>	2	0
Canarygrass	<i>Phalaris sp.</i>	1	0
Chenopodium sp.	<i>Chenopodium sp.</i>	1	0
Legume family cf.	<i>Fabaceae</i>	1	0
Nightshade	<i>Solanum sp.</i>	1	0
St. Johnswort	<i>Hypericum sp.</i>	4	0
Violet	<i>Viola sp.</i>	1	0
Wild cucumber	<i>Marah macrocarpus</i>	145	0.34
Wild cucumber cf.	<i>Marah macrocarpus</i>	17	0.03
Unidentified		232	0.55
Unidentified seed		9	0

References Cited

- Martin, A. C., and W. D. Barkley
 1961 *Seed Identification Manual*. University of California Press, Berkeley.
- Miksicek, Charles H.
 1987 Formation Processes of the Archaeobotanical Record. In *Advances in Archaeological Method and Theory*, Vol. 10, ed. by M. Schiffer, pp. 211–247. Academic Press, New York.
- Minnis, Paul E.
 1981 Seeds in Archaeological Sites: Sources and Some Interpretive Problems. *American Antiquity* 46(1):143–152.
- Minnis, Paul E. (editor)
 2004 *People and Plants in Ancient Western North America*. Smithsonian Books, Washington.

Scarry, C. Margaret

1986 Change in Plant Procurement and Production during the Emergence of the Moundville Chiefdom. Unpublished Ph.D. dissertation, Department of Anthropology, University of Michigan, Ann Arbor.

Timbrook, Jan

2007 *Chumash Ethnobotany: Plant Knowledge Among the Chumash People of Southern California*. Santa Barbara Museum of Natural History, Santa Barbara & Heyday Books, Berkeley.

Yarnell, Richard A.

1982 Problems of interpretation of archaeological plant remains of the eastern woodlands. *Southeastern Archaeology* 1(1):1-7.